

University College London
Department of Computer Science

Cryptanalysis Lab 2

J. P. Bootle

Cyclic Groups

Click on the “Ans” button to get a hint.

Shift-click on “Ans” buttons that have a green boundary to get a full solution. Click on the green square to go back to the questions.

Quiz

1. How many elements in $(\mathbb{Z}/11\mathbb{Z})^*$?

=

2. Find a single element that generates $(\mathbb{Z}/11\mathbb{Z})^*$.

=

3. What is the order of 5 in $(\mathbb{Z}/11\mathbb{Z})^*$?

=

EXERCISE 1.

(a) Let p be a prime such that $p = 2q + 1$, where q is also prime. We call p with this property a ‘strong’ prime or ‘safe’ prime. Let g



Back

be a generator of $(\mathbb{Z}/p\mathbb{Z})^*$. How can we generate a group of order q ?

Cyclic Groups in SAGE

Try out the following sequence of SAGE commands, and verify that the first 3 results match with your answers to the first 3 questions.

EXERCISE 2.

- (a) `euler_phi(11)`
- (b) `primitive_root(11)`
- (c) To find the order of 5:
 - `R = Integers(11)`
 - `a = R(5)`
 - `a.multiplicative_order()`
- (d) Compute (easy) discrete logarithms:
 - `R = Integers(11)`
 - `a = R(5)`
 - `b = a*a*a*a`
 - `a.log(b)`



(e) Compute modular square roots:

$$R = \text{Integers}(7)$$

$$a = R(3)$$

$$b = a*a$$

$$\text{mod}(2,7).\text{sqrt}()$$

The Fermat Factorisation Algorithm

Click on the green letter before each question to get a full solution.
Click on the green square to go back to the questions.

EXERCISE 3.

- (a) Given that $1309 = 47^2 - 30^2$, what is the prime factorisation of 1309?
- (b) Let N, a, b be odd, positive integers such that $N = ab$. Show that N can be expressed as the difference between two square numbers.
- (c) The incomplete function ‘Fermat’ implements a factorisation algorithm. The function takes input N , and should output a, b such



Back

that $N = ab$. Please fill in the question marks to obtain a complete implementation of the Fermat factorisation algorithm.

```
def fermat(N):
    n = ceil(sqrt(N))
    while ???:
        M = n*n-N
        m = floor(sqrt(M))
        if m == sqrt(M):
            return ???
        n = n+1
```

- (d) Use your completed code to find the factors of $N = 1488391, 1467181, 1456043$. Can you see a connection between the running time of your code and the prime factors of N ?

Rabin Cryptosystem

Click on the green letter in front of each sub-question (e.g. (a)) to see a solution. Click on the green square at the end of the solution to go back to the questions.



Back

EXERCISE 4. Let p, q be two large primes which are congruent to 3 modulo 4. Set $N = pq$.

- (a) Let $c \equiv m^2 \in \mathbb{Z}/p\mathbb{Z}$. Set $m' \equiv c^{(p+1)/4} \pmod{p}$. What is $(m')^2$?
- (b) The Rabin cryptosystem encrypts a message $m \pmod{N}$ by setting $c \equiv m^2 \pmod{N}$. Suppose that you know p, q . Use the first part of the question to describe how to decrypt a message. Hint: use the Chinese Remainder Theorem.
- (c) With a partner, generate two primes which are suitable for the Rabin cryptosystem. Now, using SAGE, write programs which can encrypt and decrypt a message. The CRT command is very useful for this.



Solutions to Exercises

Exercise 1(a) The order of g is $\phi(p) = p - 1 = 2q$. We can compute $g^2 \pmod p$, and this element will have order q , generating a subgroup of size q . \square



Exercise 3(a) We have $1309 = (47 + 30)(47 - 30) = 77 \cdot 17$.



Back

Exercise 3(b) Write $N = \left(\frac{a+b}{2}\right)^2 - \left(\frac{a-b}{2}\right)^2$. Each bracketed expression is a whole number, because N is odd, so a, b are both odd, and therefore $a \pm b$ is even. \square



Exercise 3(c) The following code implements the Fermat Factorisation algorithm.

```
def fermat(N):  
    n = ceil(sqrt(N))  
    while True:  
        M = n*n-N  
        m = floor(sqrt(M))  
        if m == sqrt(M):  
            return [n+m,n-m]  
        n = n+1
```



Exercise 3(d) The Fermat factorisation method finds factors of N as $n + m$ and $n - m$, where $N = n^2 - m^2$. The value of $n + m$ is at least \sqrt{N} and increases as n is incremented. Therefore, Fermat factorisation runs fastest on integers N which have factors close to \sqrt{N} . \square



Exercise 4(a) By Fermat's Little Theorem, we have that $(m')^2 \equiv c \pmod{p}$. □



Exercise 4(b) We can compute $c_p \equiv c \pmod{p}$ and $c_q \equiv c \pmod{q}$. Using the first part of the question, we can compute the square roots m_p with $m_p^2 \equiv c_p \pmod{p}$ and $m_q^2 \equiv c_q \pmod{q}$. Finally, we can use the Chinese Remainder Theorem to compute $m \pmod{N}$ from $m_p \pmod{p}$ and $m_q \pmod{q}$. \square



Solutions to Quizzes

Solution to Quiz: The number of elements in $(\mathbb{Z}/N\mathbb{Z})^*$ is $\phi(N)$, so in this case, the answer is $\phi(11) = 10$. ■



Back

Solution to Quiz: If we compute the powers of 2 modulo 11, we get 2, 4, 8, 5, 10, 9, 7, 3, 6, 1, so 2 is a generator. Alternatively, by Lagrange's Theorem, the order of an element divides the size of the group. The size of the group is 10, so the only possibilities for the order of an element are 1, 2, 5, and 10. A group generator should have order 10 to generate every group element, so to check that 2 is a generator, we just have to check that $2^2 \neq 1 \pmod{11}$, and $2^5 \neq 1 \pmod{11}$, implying that 2 has order 10. ■



Solution to Quiz: The smallest n such that $5^n = 1 \pmod{11}$ is $n = 5$. Alternatively, by Lagrange's Theorem, the order of an element divides the size of the group. The size of the group is 10, so the only possibilities for the order of an element are 1, 2, 5, and 10. Therefore, it is enough to check that $5^2 \not\equiv 1 \pmod{11}$, and $5^5 = 1 \pmod{11}$. ■

